

Solitonic photons and intermediate vector bosons

M. Temple-Raston *

Department of Mathematics and Statistics,
Concordia University,
Montréal, Québec, Canada

February 1, 2008

Abstract

A four-dimensional topological field theory is introduced which generalises $B \wedge F$ theory to give a Bogomol'nyi structure. A class of non-singular, finite-action, stable solutions to the variational field equations is identified. The solitonic solutions are analogous to the instanton in Yang-Mills theory. The solutions to the Bogomol'nyi equations in the topologically least complicated $U(1)$ theory have a covariant phase space of dimension four—the same as that for photons. The dimensional reduction of the four-dimensional Lagrangian is also examined. Bogomol'nyi $U(2)$ solitons resembling the intermediate vector bosons Z_0 , $W^\pm$ are identified.

1 Introduction

The photon is a non-singular, finite-energy field configuration with a definite frequency ν , wave vector $\mathbf{k}$, and helicity ± 1 . Furthermore, the energy of the photon, E , and its momentum, $\mathbf{p}$, satisfy the dispersion relation $E = c |\mathbf{p}|$.

*current address: Centre for Industrial Control, Mechanical Engineering, Concordia University, Montreal, Quebec, Canada

The dispersion relation implies that the phase space, $\mathcal{P}$, exists and must presumably be a well-behaved manifold (e.g., Hausdorff). If the photon is solitonic (in the general sense of the word), then the phase space becomes the ‘covariant’ phase space. The covariant phase space is given by the moduli space of solutions to the photonic time-dependent field equations. At present, field equations for a solitonic photon are unknown. However, it is a strong requirement on the possible field equations to have a well-behaved covariant phase space, $\mathcal{P}$. If additionally we require that the photonic covariant phase space be finite dimensional and in particular of dimension four, we now possess very strong conditions on classical field theories which dictate the terms under which a theory is acceptable. Subject to these physical requirements on the covariant phase space, we introduce in this paper a Lagrangian field theory which possesses Bogomol’nyi solitons with properties similar to the photon. The theory also contains stationary solitons that resemble other force mediating particles. They occur as Bogomol’nyi solitons to the time-reduced field equations.

We now summarize the sections of this paper. In the next section we introduce two classical, generally covariant Lagrangian field theories defined on a four-manifold. The natural Bogomol’nyi structures are identified. Non-singular solutions to the Bogomol’nyi equations are shown to be stable under small perturbations. When the theory is defined on a compact, complex Kähler surface a finite dimensional covariant phase space is found for the Bogomol’nyi solitons, with good topological behaviour. The covariant phase space for abelian (diagonal) Bogomol’nyi solitons over various compact, complex surfaces is examined in detail. In section three we dimensionally reduce the four-dimensional topological field theory using a time-symmetry to a three-dimensional topological field theory. The three-dimensional field theory permits electric, magnetic and dyonic monopoles. From the general arguments in [11] the particle spectrum and classical mass compare favourably with that of the intermediate vector bosons.

2 Topological solitons

Let $\pi : P \rightarrow M$ be a principal G -bundle over an oriented, compact, connected four manifold M , and denote by $E = P \times_G \mathcal{G}$, the associated adjoint vector bundle where $\mathcal{G}$ is the Lie algebra of G . Denote by $\mathcal{A}(P)$ the space of

connections on P . Let $A, B \in \mathcal{A}(P)$. We introduce local coordinate indices $\mu = 0, \dots, 3$ on M . The $\mathcal{G}$ -valued connections or vector potentials, A_μ and B_μ , induce exterior covariant derivatives $D_\mu^A = \partial_\mu + A_\mu$ and $D_\mu^B = \partial_\mu + B_\mu$ on the associated adjoint vector bundle E . The curvatures H and K are defined by $2D_{[\mu}^A D_{\nu]}^A s = sH_{\mu\nu}$ and $2D_{[\mu}^B D_{\nu]}^B s = sK_{\mu\nu}$. As a result, H and K are two-forms on M taking values in E . We introduce a Lagrangian Action functional constructed in such a way to produce invariants and to be geometrically meaningful. Physically, the invariants and geometrical interpretation allow us to argue the existence and stability of our solitonic solutions. The Lagrangian theory is given by the functional

$$\begin{aligned} \mathcal{L}_\pm(A, B) = & \int_M H_{[\mu\nu]}^a K_{\lambda\rho]}^b \operatorname{tr}(T^a) \operatorname{tr}(T^b) d^4x \\ & \pm \frac{1}{2} \int_M K_{[\mu\nu]}^a K_{\lambda\rho]}^b \operatorname{tr}(T^a T^b) d^4x. \end{aligned} \quad (1)$$

The generators of the Lie algebra are denoted by T^a . For gauge groups where $\operatorname{tr}(T^a) = 0$, the Lagrangian reduces to the second integral—these are the topological field theories studied by Baulieu and Singer [2]. We do not a priori restrict the Lie group.

The variational field equations are

$$D_{[\mu}^A K_{\nu\rho]} = 0, \quad D_{[\mu}^B H_{\nu\rho]} = 0. \quad (2)$$

The Lagrangian can be rewritten as

$$\begin{aligned} 2\mathcal{L}_\pm = & \pm \int_M < (H_{[\mu\nu]} \otimes I_E \pm I_E \otimes K_{[\mu\nu]}) (H_{\lambda\rho]} \otimes I_E \pm I_E \otimes K_{\lambda\rho]} > d^4x \\ & \mp \int_M H_{[\mu\nu]}^a H_{\lambda\rho]}^b \operatorname{tr}(T^a T^b) d^4x. \end{aligned} \quad (3)$$

I_E denotes the identity transformation on the adjoint bundle, E . The Lagrangian in (3) is now in Bogomol'nyi form. We need to interpret the first term in $\mathcal{L}_\pm$. Let E_A and E_B be the adjoint vector bundles equipped with D_μ^A or D_μ^B , respectively. E_B^* is the dual bundle to E_B . Now, recall that the curvature of the tensor product bundle $E_A \otimes E_B^*$ is given by $(\Omega_{E_A \otimes E_B^*})_{\mu\nu} = H_{\mu\nu} \otimes I_E - I_E \otimes K_{\mu\nu}$ [10]. Likewise, the curvature of $E_A \otimes E_B$ is given by $(\Omega_{E_A \otimes E_B})_{\mu\nu} = H_{\mu\nu} \otimes I_E + I_E \otimes K_{\mu\nu}$. Since the trace inner product on E generalizes naturally to an inner product on $E \otimes E$ and $E \otimes E^*$, we have signified both inner products by $< >$ in (3). The appropriate inner product is determined by the context. Under an infinitesimal perturbation of the vector potential, the integrand in the first term of both Lagrangians (3) changes by

an exact form. The exact form vanishes when integrated over compact M , so that the first term in (3) is invariant under small perturbations.

In our view, the more interesting theory is that of the Lagrangian $\mathcal{L}_-$, so we concentrate on it. The Bogomol'nyi equations in (3) for $\mathcal{L}_-$ are seen to be a vanishing curvature condition on the tensor product bundle $E_A \otimes E_B^*$:

$$(\Omega_{E_A \otimes E_B^*})_{\mu\nu} = H_{\mu\nu} \otimes I_E - I_E \otimes K_{\mu\nu} = 0. \quad (4)$$

Solutions to (4) automatically satisfy the variational field equations (2). The equations (4) give useful information about H and K ; they imply that H and K are projectively flat,

$$H_{\mu\nu} = K_{\mu\nu} = iF_{\mu\nu}I, \quad (5)$$

where F is a real-valued two form on M . The projective flatness reduces the gauge group down to $U(n)$. Non-trivial solitonic solutions (A_μ, B_μ) to the Bogomol'nyi equations (4) are stable under a small perturbation when the Lagrangian Action is invariant and

$$\int_M \text{tr}(H_{[\mu\nu}H_{\lambda\rho]}) d^4x \quad (6)$$

is non-vanishing.

There are clear similarities between what we have constructed above and the (anti-)self-dual instanton in Yang-Mills theory. We, however, will not call these solutions ‘instantons’. Because physicists have come to associate instantons with Euclidean four-space—only trivial instanton solutions exist in Minkowski space. Unlike instantons, the Bogomol'nyi solutions in these topological field theories are not affected by the metric signature of space-time. In light of this, we call non-trivial, stable, non-singular solutions to the Bogomol'nyi equations (5) topological solitons. The term ‘soliton’ is used rather generally in this paper, and does not suggest that the solutions are connected to inverse scattering or that soliton-soliton interactions are trivial. Note that a space-time metric is not needed to define the Lagrangian field theory (1). Therefore the underlying manifold M need not be space-time at all. In view of this, we define M to be a four-dimensional manifold which defines the *topology* of the field configuration on space-time. The topology is defined by the number of isolated zeros and the multiplicities in the field configuration [3]. The topology of M , call it the topological field manifold,

defines a set of topologically equivalent field configurations on space-time. If M is indeed endowed with the space-time metric and viewed as space-time, then this reduces to a particular example of the general theory we propose.

We now restrict to those topological solitons (A_μ, B_μ) of the Bogomol'nyi equations (4) for which $A_\mu = B_\mu$. Let us call these solutions ‘diagonal topological solitons’. The variational field equations (2) become the Bianchi identities. The only other condition on the vector potential, A_μ , comes from the Bogomol'nyi equations, which require that A_μ be projectively flat. We must now state our connection to Maxwell's theory of electromagnetism. Consider first a $U(1)$ gauge theory. Since in a $U(1)$ gauge theory projective flatness is a vacuous requirement, the abelian vector potential is completely unconstrained. We are led therefore to introduce Maxwell's equations. We do this by requiring that $F_{\mu\nu}$ in equation (5) is a Maxwell field (Faraday tensor). The Bianchi identities for H and K imply automatically that $dF = 0$. The second half of the Maxwell equations, $*d*F = j$, however, we must introduce by hand. The introduction of Maxwell's equations at the level of $F_{\mu\nu}$ implies that in $U(n)$, $n > 1$, gauge theories $H_{\mu\nu} = iF_{\mu\nu}I$ satisfies the non-homogeneous Yang-Mills equations, $*D^A *H = J \equiv jI$. In the following, we permit the second half of the Maxwell equations be relaxed if needed.

So far, the Lagrangian introduced above has yielded a Bogomol'nyi structure. The Bogomol'nyi structure suggests that interesting particle-like solutions to the theory might be found. However, we have also seen that in the case of the abelian gauge group only vacuous conditions remain in the theory. Nothing further remains in our theory to determine these solitons. The way forward is made clearer by the non-abelian theory. The Bogomol'nyi equations (5) are non-trivial in the non-abelian theory; they require that the vector potential be projectively flat. Projectively flat vector potentials are closely related to Einstein-Hermitian connections, which in turn express in differential geometry the algebraic geometric concept of ‘stable vector bundle’ [10]. As a result of this close connection to stable vector bundle, projectively flat vector potentials often have a well-behaved moduli space—like self-dual instantons and BPS magnetic monopoles. In the four-dimensional theory studied in this section, the moduli space of topological solitons is equivalent to the covariant phase space (or, the space of motions).

To establish the link between projectively flat and Einstein-Hermitian connections, we must introduce further geometrical structure into the theory. We require that the topological field manifold M be a compact, complex

Kähler surface. Clearly we have introduced the Kähler structure (g, Φ) on M at the expense of the space-time structure. M is no longer space-time. The Kähler metric and Kähler form are given by g and Φ , respectively. We also restrict the $U(1)$ theory, which is all that interests us in this section, to holomorphic Hermitian line bundles $(L, \bar{\partial}, h = \langle \rangle)$ over M . The set of connections compatible with the added geometric structure (the Kähler metric g , the complex structure $\bar{\partial}$) are denoted by $\mathcal{D}^{1,1}(L, h)$. Einstein-Hermitian connections are defined as the connections with constant mean curvature, $K(A) = c$, compatible with the holomorphic Hermitian structure. Let $\mathcal{E}(L, h)$ denote the set of all Einstein-Hermitian connections in $\mathcal{D}^{1,1}(L, h)$. Kobayashi has shown that all connections in $\mathcal{D}^{1,1}(L, h)$ are Einstein-Hermitian up to a conformal transformation of the Hermitian structure, $h[9]$. The constant c is given by

$$c = \frac{2\pi}{V} \deg(L),$$

where $V = \text{Vol}(M)$ and $\deg(L) = \int_M c_1(L) \wedge \Phi$. c is a topological invariant when M is Kähler (when Φ is closed), and depends only on the cohomology classes of Φ and $c_1(L)$ [10].

As mentioned before, the moduli space of Einstein-Hermitian connections is equivalent to the covariant phase space for the topological solitons defined by the Bogomol'nyi equations (5) on (M, g, Φ) . The covariant phase space has been studied extensively by H.J. Kim [8]. Kim proves that the covariant phase space $\mathcal{M} = E(L, h)/U(1)$ is a non singular Kähler manifold if $H^2(M, \text{End}^0(L)) = 0$, and

$$\deg(M) \equiv \int_M c_1(M) \wedge \Phi \geq 0. \quad (7)$$

When the inequality (7) is satisfied, the complex dimension of the covariant phase space of abelian diagonal topological solitons is found to be

$$\dim_{\mathbf{C}}(M) = q, \quad (8)$$

where $q \equiv h^{(0,1)}$ is the irregularity of M . The irregularity is one of the Hodge numbers in complex cohomology theory, defined generally by $h^{(p,q)} \equiv \dim_{\mathbf{C}} H^{p,q}(M)$. The natural symplectic structure on the covariant phase space is inherited from the Kähler structure on the configuration manifold, M :

$$\Theta(a, b) = \int_M \text{tr}(a \wedge b) \wedge \Phi,$$

where $a, b \in T\mathcal{M}$ [12].

Which field configuration topologies lead to interesting covariant phase spaces in this theory? The simplest topology, that of the K3 surface, has irregularity $q = 0$. The dimension of the covariant phase space with this field topology is therefore zero, from equation (8). This phase space is too small to be interesting. The next simplest field configuration, that of the flat complex two-torus, $T^2 = \mathbf{C}^2/\Gamma$, has irregularity $q = 2$. For this field configuration the real dimension of the covariant phase space is four. While the dimension of the covariant phase space is non-zero, the stability of the topological solitons under a small perturbation is still under question. From (6), we recall that the stability for vector bundles of rank $r > 1$ is assured when the second Chern class is non-zero. For line bundles, however, equation (6) is not generally a topological invariant. We can make (6) a topological invariant by interpreting $F_{\mu\nu}$ in the Bogomol'nyi equation (14) as a curvature on the tangent bundle, TM . Equation (6) is then proportional to the signature of the underlying four-manifold, M . Thus, the stability of the topological soliton in the abelian theory is argued strictly from the topology of the topological field manifold, M . Returning to the flat complex two-torus, recall that for T^2 the irregularity (q) and the geometric genus (p_g) take the values $q = 2$ and $p_g = 1$. Also $c_1^2(T^2) = 0$. From the Riemann-Roch formula

$$c_1^2(M) + c_2(M) = 12(1 - q + p_g)$$

we conclude that the signature vanishes for T^2 . Therefore solutions to the Bogomol'nyi equations with a topological field manifold $M = T^2$ are unstable. The next topological field manifold we examine is found in the Enriques-Kodaira classification of compact, complex surfaces (free from exceptional curves). There are two possibilities: minimal elliptic surfaces and minimal algebraic surfaces of 'general type'. We examine here the simpler of the two. Let M be a minimal elliptic ($c_1^2(M) = 0$) surface with irregularity $q = 2$. Again, the covariant phase space is of real dimension four. But in addition now, stability is assured for any topological field manifold M with geometric genus $p_g > 1$. Thus we have shown that the real dimension of the covariant phase space for $U(1)$ diagonal topological solitons (understood to be stable) on minimal elliptic complex surfaces with geometric genus greater than one, is four. Recall that the photon has a phase space (space of motions) of real dimension four as well.

The explicit construction of topological solitons on minimal elliptic topological field manifolds, M , is probably a difficult problem—as is the case here, it is often much easier to construct the moduli space. A connection between the (diagonal) topological soliton and the photon is further suggested by a natural mechanism by which this theory has no anti-solitons, in agreement with the notable absence of the anti-photon in nature. This can be achieved if the transformations for time and parity reversal of H and K are:

$$\begin{aligned} T : H &\longmapsto H & P : H &\longmapsto -H \\ T : K &\longmapsto -K & P : K &\longmapsto -K \end{aligned}$$

Time reversal takes the Lagrangian Action $\mathcal{L}_-$ to $-\mathcal{L}_+$ in equations (3). The Bogomol'nyi equations for L_+ are similar in appearance to those for L_- :

$$(\Omega_{E_A \otimes E_B^*})_{\mu\nu} = H_{\mu\nu} \otimes I_E + I_E \otimes K_{\mu\nu} = 0. \quad (9)$$

An index computation for these Bogomol'nyi equations yields the equations

$$H_{\mu\nu} = -K_{\mu\nu} = iF_{\mu\nu}I.$$

It is clear that diagonal topological solitons are no longer diagonal after time reversal. On the other hand, under transformations of the parity the Lagrangian Action is the same up to a overall sign—the mirror image of the diagonal topological soliton is again a solution to the Bogomol'nyi equations. This is compatible with two helicity states.

3 Topological vector bosons

In the previous section we studied the abelian topological field theory defined by equation (1). We found a Bogomol'nyi structure with topological solitons that resemble photons. In this section we examine the non-abelian theory. If indeed the topological solitons of section two can be viewed as photons, then it is natural to expect other intermediate vector bosons to be found in a similar manner. Our interest in this section now lies in topological solitons with rest mass. Without loss of generality, we can restrict our investigation to stationary topological solitons. At this point one is reminded of the Montonen-Olive conjecture, which argues that intermediate

vector bosons should arise as solitons in a Lagrangian field theory, analogous to the way that BPS magnetic monopoles are found in Yang-Mills-Higgs theory [11]. Since BPS magnetic monopoles and instantons are related by a gauge symmetry, it is to be expected that we try the same trick here on the topological solitons found in the previous section. By assuming a gauge symmetry in time, the dimensionally reduced Lagrangian Action gives the energy functional $\mathcal{E}(A, B, \Phi_A, \Phi_B)$ equal to

$$\begin{aligned} \int_{M_3} K_{[\mu\nu]}^a (D_{[\rho]}^B \Phi_B)^b \text{tr}(T^a T^b) &- K_{[\mu\nu]}^a (D_{[\rho]}^A \Phi_A)^b \text{tr}(T^a) \text{tr}(T^b) \\ - H_{[\mu\nu]}^a (D_{[\rho]}^B \Phi_B)^b \text{tr}(T^a) \text{tr}(T^b) &+ (A \leftrightarrow B, \Phi_A \leftrightarrow \Phi_B). \end{aligned} \quad (10)$$

We have followed the procedure of dimensional reduction in [4]. Note that in (10) there are two curvatures H, K and two Higgs fields Φ_A, Φ_B corresponding to the two connections A, B . We have added by hand the last term in (10) to symmetrise the energy functional in the dependent fields. The symmetrisation although probably desirable in principle in section two, made no change to the development of diagonal topological solitons in the four-dimensional theory, so we left it out. The energy functional looks rather complicated, but it does possess analogous geometric structures to those found in the four dimensional theory. The variational field equations arising from (10) are

$$\begin{aligned} D^B H &= 0, & D^B D^A \Phi_A &= [H, \Phi_B], \\ D^A K &= 0, & D^A D^B \Phi_B &= [K, \Phi_A]. \end{aligned} \quad (11)$$

The energy functional (10) can be rewritten as

$$\begin{aligned} \mathcal{E} = \int_{M_3} &< (H_{[\mu\nu]} \otimes I_E - I_E \otimes K_{[\mu\nu]}) \wedge (D_{[\rho]}^A \Phi_A \otimes I_E - I_E \otimes D_{[\rho]}^B \Phi_B) > \\ - \int_{M_3} &< (D_{[\mu]}^A \Phi_A \otimes I_E) \wedge (H_{\nu\rho]} \otimes I_E) > + (A \leftrightarrow B, \Phi_A \leftrightarrow \Phi_B). \end{aligned} \quad (12)$$

In this equation we begin to see the geometrical structure emerge out of the energy functional (10). Let E_A and E_B be the vector bundle E equipped with either the connection A or B , respectively. Again, the curvature of the tensor product bundle $E_A \otimes E_B^*$ is given by $\Omega_{E_A \otimes E_B^*} = H_{\mu\nu} \otimes I_E - I_E \otimes K_{\mu\nu}$. This curvature expression appears in (12). By defining $\Phi \equiv \Phi_A \otimes I_E - I_E \otimes \Phi_B$, then $D_{E_A \otimes E_B^*} \Phi = D^A \Phi_A \otimes I_E - I_E \otimes D^B \Phi_B$, and the first integral in (12) is therefore a topological invariant. The Bogomol'nyi equations arising from equation (10) are

$$\begin{aligned} H_{\mu\nu} \otimes I_E &= I_E \otimes K_{\mu\nu} \\ D_\mu^A \Phi_A \otimes I_E &= I_E \otimes D_\mu^B \Phi_B \end{aligned} \quad (13)$$

The first equation in (13) is a zero curvature condition on the tensor product bundle $E_A \otimes E_B^*$. From an indices computation we conclude that

$$\begin{aligned} H_{\mu\nu} &= K_{\mu\nu} = iF_{\mu\nu}I_E, \\ D_\mu^A \Phi_A &= D_\mu^B \Phi_B = iE_\mu I_E. \end{aligned} \quad (14)$$

F and E are a real-valued two-form and one-form on M_3 , respectively. Solutions to (14) automatically satisfy the variational field equations (11). Recalling section two, solutions to the first equation are projectively flat connections [10]. For line bundles this carries no extra information, but for bundles of rank greater than one projective flatness is a strong condition. Unlike the theory of BPS magnetic monopoles, however, the energy functional (12) is saturated at the Bogomol'nyi energy with either Bogomol'nyi equation in (14) satisfied. Of course the field configurations must still satisfy the second-order variational field equations. The Bogomol'nyi energy is given by

$$\mathcal{E} = - \int_{M_3} (D_{[\rho}^A \Phi_A)^a H_{\mu\nu]}^b \text{tr}(T^a T^b) - \int_{M_3} (D_{[\rho}^B \Phi_B)^a K_{\mu\nu]}^b \text{tr}(T^a T^b). \quad (15)$$

The added flexibility in obtaining the Bogomol'nyi energy, a feature not present in the theory of magnetic monopoles, will lead to electric monopoles.

Since topological monopoles, if they exist, are analogous to the BPS magnetic monopole field configurations, we shall use the same symmetry breaking mechanism [5]. We place the solitonic core region at the origin. Let G and G_o be compact and connected gauge groups, where the group G_o is assumed to be embedded in G . The gauge group of the core region G is spontaneously broken to G_o outside of the core region when the Higgs field is covariantly constant, $D\Phi = 0$. In regions far from the core ($r \rightarrow \infty$) where we assume that $D^A \Phi_A = 0$, it can be shown that

$$H = \Phi_A F_A, \quad (16)$$

where $F_A \in \Lambda^2(M_3, E_{G_o})$, a two-form on M_3 taking values in the G_o -Lie algebra bundle, denoted by E_{G_o} here [5]. An equivalent expression to (16) can be written when $D^B \Phi_B = 0$. We adopt units so that $\Phi^2 = 1$ when $r \gg 1$ and where spontaneous symmetry breaking has occurred. When $G = U(n)$ and $G_o = U(1)$, F_A becomes a pure imaginary two-form on M_3 .

The Bogomol'nyi solitons defined by (14) have an energy coming from (15) topologically fixed by

$$\begin{aligned}\mathcal{E} &= -\int_{M_3} d(\text{tr}(\Phi_A H)) - \int_{M_3} d(\text{tr}(\Phi_B K)) \\ &= -\int_{S^2} \text{tr}(\Phi_A H) - \int_{S^2} \text{tr}(\Phi_B K)\end{aligned}\tag{17}$$

where S^2 is a large sphere surrounding the monopole core. Details of the solitonic particles in this theory now depend on the extent to which they satisfy the Bogomol'nyi equations (14) and are 'spontaneously broken'.

Consider first non-singular particle-like solutions to *both* Bogomol'nyi equations in (14). Substituting (16) into (17) and using the normalization condition $\Phi^2 = 1$ for both Higgs fields, the energy is fixed by $-\int F_A - \int F_B$. Let us conventionally interpret $\int F_A/2\pi$ as the magnetic charge (g), and $\int F_B/2\pi$ as the electric charge (q). We can view F_A and F_B as curvatures on the line bundles L_A and L_B determined by Φ_A and Φ_B ; from (16) F_A and F_B are the projections of H and K on L_A and L_B . The magnetic and electric charges are thereby the Chern numbers associated to the complex line bundles with curvatures F_A and F_B , respectively, and in this theory are thereby quantized at the classical level. The stability of the solitonic particle is argued from the topological interpretation, a winding number, that can be given to (17). When both equations in (13) are satisfied the topological stability is assured if either the electric or magnetic charge is non-vanishing. To this point, we have used only half of the Bogomol'nyi equations, $D^A \Phi_A = D^B \Phi_B = 0$. From the projective flatness of the curvatures in the Bogomol'nyi equations (14), $H = K = FI_E$, and (16) we conclude that $F = \varphi_A F_A = \varphi_B F_B$ where $\Phi_A = \varphi_A I_E$ and $\Phi_B = \varphi_B I_E$. From this we find that

$$\int_{S^2} F_A = \int_{S^2} F_B.\tag{18}$$

Therefore non-singular, stable, particle-like solutions to the Bogomol'nyi equations (14) are dyons. To obtain electric monopoles there would appear to be two possibilities, both resulting from a weakening of the Bogomol'nyi equations (14).

In the first possibility, one or the other of the non-abelian gauge fields, H or K , does not break to $U(1)$ in the far-field, so that non-abelian gauge field passes unnoticed through the detector. This corresponds to the case where, for example, $H^A = K^B = iFI_E = F_B \Phi_B$, but asymptotically $D^B \Phi_B = 0$,

$D^A\Phi_A \neq 0$. The Bogomol'nyi bound (15) in this case becomes

$$\mathcal{E} = - \int_{S^2} F_B \operatorname{tr}(\Phi_A) \operatorname{tr}(\Phi_B) - \int_{S^2} F_B. \quad (19)$$

The second term is -2π times the electric charge. Since the projective flatness reduces the gauge group to $U(n)$, we have used the Killing-Cartan form for the bundle inner product, $\langle \cdot, \cdot \rangle$. The first integral in (19) vanishes when Φ_A is traceless. An explicit example of an electric monopole is $A = B$ projectively flat, and Φ_A, Φ_B are any sufficiently differentiable functions on M taking values in $SU(n)$ and $U(n)$, respectively, with Φ_B covariantly constant asymptotically ($D^B\Phi_B = 0$ for $r \rightarrow \infty$). The full variational field equations are satisfied.

The second possibility allows both gauge fields to break far from the core using $D^A\Phi_A = D^B\Phi_B = 0$, but the two gauge far-fields $H^A = F_A\Phi_A$ and $K^B = F_B\Phi_B$ are completely decoupled so that F_A and F_B and their respective topological charges are unrelated. That is, the projective flatness in the Bogomol'nyi equations is relaxed. The electric and magnetic charges in this case become, respectively,

$$2\pi q = - \int_{S^2} F_B, \quad 2\pi g = - \int_{S^2} F_A,$$

with no relation between q and g analogous to (18).

In this section we introduced a topological field theory with a Bogomol'nyi structure permitting BPS electric, magnetic and dyonic monopoles. From the general arguments in [11] the particle spectrum and classical mass compare favourably with that of the intermediate vector bosons. Using the general relationship between supersymmetry and the Bogomol'nyi equations it can be shown that there are no quantum corrections to the classical mass spectrum [14][6]. For $G = U(2)$, which is double covered by $SU(2) \times U(1)$, the solitons are associated with the $W^\pm$ and Z_0 intermediate vector bosons. In this theory $W^\pm$ have energies of opposite sign so that, presumably, one is the anti-particle of the other, as required. The Z_0 probably corresponds to the case where $H^A = K^B = iFI_E$, but $D^A\Phi_A \neq 0$ and $D^B\Phi_B \neq 0$. The gauge far-fields for the Z_0 are thereby non-abelian and pass unnoticed through conventional detectors, and have a stabilizing energy given by

$$\mathcal{E} = - \int_{S^2} F \operatorname{tr}(\Phi_A) - \int_{S^2} F \operatorname{tr}(\Phi_B)$$

This completes the work begun in [13].

4 Conclusion

We have introduced a Lagrangian field theory which contains solitonic particles and possesses a well-behaved covariant phase space, relating the topology of the field configuration to the dimension of the covariant phase space. It was suggested that these classical field configurations provide a provisional basis from which a realistic solitonic model for force mediating bosons could be found. We have not addressed in this paper the most important question: interaction dynamics (soliton-soliton interactions and soliton-matter interactions).

References

- [1] M. Atiyah, N. Hitchin, *The Geometry and Dynamics of Magnetic Monopoles* (Princeton U.P., New Jersey, 1988).
- [2] J-M. Baulieu, I. Singer, *Proceedings of conformal field theory and related topics* (Annecy, France, March 1988).
- [3] S-S Chern, *Complex Manifolds Without Potential Theory*, 2nd ed. (Springer-Verlag, New York, 1979).
- [4] P. Forgács, N.S. Manton, *Comm. Math. Phys.* **72** (1980) 15.
- [5] P. Goddard, D. Olive, *Rep. Prog. Phys.* **41** (1978) 1357.
- [6] Z. Hlousek, D. Spector, *Nucl. Phys.* **B397** (1993) 173.
- [7] G.T. Horowitz, *Comm. Math. Phys.* **125** (1989) 46.
- [8] H.J. Kim, *Math. Z.* **195** (1987) 143.
- [9] S. Kobayashi, S., in *Global Riemannian Geometry*, eds. T.J. Willmore and N. Hitchin (Ellis Horwood, Chichester, 1984) 60.
- [10] S. Kobayashi, *Differential Geometry of Complex Vector Bundles* (Princeton U.P., New Jersey, 1987).
- [11] C. Montonen, D.I. Olive, *Phys. Lett.* **72B** (1977) 117.

- [12] S. Mukai, S., *Inven. Math.* **77** (1984) 101.
- [13] M. Temple-Raston, *Lett. Math. Phys.* **23** (1991) 91.
- [14] E. Witten, D. Olive, *Phys. Lett.* **78B** (1978) 97.